

Lesson 8

Carbon Through the Seasons



Carbon Through the Seasons

NYS Intermediate Level Science

Standard 1: Analysis, Inquiry and Design/Scientific Inquiry

- S3.1a Organize results, using appropriate graphs.
- S3.2h Use and interpret graphs and data tables.

Standard 6: Interconnectedness

- 5.2 Observe patterns of change in trends or cycles and make predictions on what might happen in the future.

Standard 4: The Living Environment

- 6.2b The major source of atmospheric oxygen is photosynthesis. Carbon dioxide is removed from the atmosphere and oxygen is released during photosynthesis.
- 6.2c Green plants are the producers of food which is used directly or indirectly by consumers.
- 7.2d Since the Industrial Revolution, human activities have resulted in major pollution of air, water, and soil. Pollution has cumulative ecological effects such as acid rain, global warming, or ozone depletion. The survival of living things on our planet depends on the conservation and protection of Earth's resources.

Standard 4: The Physical Setting

- 2.2j Climate is the characteristic weather that prevails from season to season and year to year.
- 2.2r Substances enter the atmosphere naturally and from human activity. Some of these include carbon dioxide, methane, and water vapor. These substances can affect weather, climate, and living things.

Next Generation Science Standards

Science and Engineering Practices:

- 2. Developing and using models
- 4. Analyzing and interpreting data

Grade 6

- ESS3-5. Ask questions to clarify evidence of the factors that have caused the rise in global temperatures over the past century.

Grade 7

- ESS2-1. Develop a model to describe the cycling of Earth's materials and the flow of energy that drives this process.



Common Core State Standards

ELA in the Content Areas - Grades 6-8

CCSS.ELA-Literacy.RST.6-8.7

Integrate quantitative or technical information expressed in words in a text with a version of that information expressed visually (e.g., in a flowchart, diagram, model, graph, or table).

Common Core State Standards - Mathematics Standards for Mathematical Practice

CCSS.Math.Practice.MP2

Reason abstractly and quantitatively.

CCSS.Math.Practice.MP4

Model with mathematics.

Grade 6

CCSS.Math.Content.6.NS.C.8

Solve real-world and mathematical problems by graphing points in all four quadrants of the coordinate plane. Include use of coordinates and absolute value to find distances between points with the same first coordinate or the same second coordinate.



Carbon Through The Seasons

*Understanding the carbon cycle is essential for climate literacy. This lesson plan, **Carbon Through The Seasons** is from the US Environmental Protection Agency, and was adapted from a carbon cycle lesson developed by the National Oceanic and Atmospheric Administration (NOAA): www.esrl.noaa.gov/gsd/outreach/education/poet/CO2-Seasons.pdf*



CARBON THROUGH THE SEASONS

DESCRIPTION

In this lesson plan, students learn about the carbon cycle and understand how concentrations of carbon dioxide (CO₂) in the Earth's atmosphere vary as the seasons change. Students also learn that even with these seasonal variations, the overall amount of CO₂ is increasing in the atmosphere as a result of people's activities, which are changing the natural carbon cycle.

BACKGROUND

Carbon is a chemical element that is found all over the world and in every living thing. Oxygen is another element that's found in the air we breathe. When carbon and oxygen bond together, they form a colorless, odorless gas called CO₂. In the Earth's atmosphere, CO₂ is a greenhouse gas, which means it traps heat. This "greenhouse effect" naturally helps to keep the Earth's temperature at a level that can support life on the planet.

The atmosphere isn't the only part of the Earth that has carbon. The oceans store large amounts of carbon, and so do plants, soil, and deposits of coal, oil, and natural gas deep underground. Carbon constantly moves from one part of the Earth to another through a natural repeating pattern called the carbon cycle. The carbon cycle helps to maintain a balanced level of CO₂ in the Earth's atmosphere.

But right now, people are changing this natural balance by adding more CO₂ to the atmosphere whenever we burn fossil fuels (such as coal, oil, and natural gas)—whether it's to drive our cars, use electricity, or make products. This extra CO₂ is being added to the atmosphere faster than natural processes can remove it, causing the atmosphere to trap more heat and causing the Earth's average temperature to rise. Scientists have found that the recent levels of CO₂ in the atmosphere are abnormally high compared with the long-term historical trend, and these levels are continuing to increase at an unprecedented rate.

The amount of CO₂ found in the atmosphere varies over the course of a year. Much of this variation happens because of the role of plants in the carbon cycle. Plants use CO₂ from the atmosphere, along with sunlight and water, to make food and other substances that they need to grow. They release oxygen into the air as a byproduct. This process is called photosynthesis. Another process that is part of the carbon cycle is respiration, by which plants and animals take up oxygen and release CO₂ back into the atmosphere.

When plants are growing, photosynthesis outweighs respiration. As a result, plants take more CO₂ out of the atmosphere during the warm months when they are growing the most. This can lead to noticeably lower CO₂ concentrations in the atmosphere. Respiration occurs all the time, but dominates during the colder months of the year, resulting in higher CO₂ levels in the atmosphere during those months.



TIME:

Introduction: 60–90 minutes

LEARNING OBJECTIVES:

Students will:

- Learn about the carbon cycle
- Understand how seasonal variations affect global atmospheric CO₂ concentrations
- Understand how CO₂ concentrations in the atmosphere are changing overall in recent decades

NATIONAL SCIENCE STANDARDS:

- Content Standard A: Science as inquiry
- Content Standard D: Earth and space science
- Content Standard E: Science and technology

ADAPTED FROM:

National Oceanic and Atmospheric Administration (NOAA):

<http://www.esrl.noaa.gov/gsd/outreach/education/poet/CO2-Seasons.pdf>.



Cornell University

CARBON THROUGH THE SEASONS

Carbon Sources and Sinks

A *carbon source* is any process or activity that releases carbon into the atmosphere. Both natural processes and people's activities can be carbon sources. A *carbon sink* takes up or stores carbon on the Earth.

The Northern and Southern Hemispheres have opposite seasons. If both hemispheres had roughly the same amount of plant life, we might expect their seasonal effects on the carbon cycle to cancel each other out. However, if you look at a map, you'll see that the Northern Hemisphere has more land than the Southern Hemisphere and a lot more plant life (especially considering that Antarctica has almost none). As a result, global CO₂ concentrations show seasonal differences that are most heavily influenced by the growing season in the Northern Hemisphere.

Scientists monitor the amount of CO₂ and other gases in the atmosphere at stations such as the Mauna Loa Observatory in Hawaii.

The Mauna Loa Observatory is one of the sites that have helped scientists determine that CO₂ levels in the atmosphere have increased significantly in recent decades and that these levels are continuing to rise at a rapid rate. CO₂ stays in the atmosphere for long enough that it is able to spread fairly evenly around the world, so even measurements from a single site (like Mauna Loa) can be representative of global average CO₂ concentrations.

GROUPS

Students should work in teams (group size of five is optimal). Each group should be assigned one set of years of "Mauna Loa Observatory Data" (see worksheets; each group will receive 10 to 12 years of data), and each student should receive one copy of his/her group's assigned data.

MATERIALS

- "Mauna Loa Observatory Data:" assign one time period per group, and give a copy of the data for the selected time period to each student in the group.
- A copy of the "Mauna Loa Worksheet" for each group
- A copy of the "Mauna Loa Monthly Average CO₂ Concentrations, 1958 to 2011" graph for each group
- Calculators
- Graph paper for each group
- Colored pencils



CARBON THROUGH THE SEASONS

VOCABULARY

Carbon:

A chemical element that is essential to all living things. Carbon combines with other elements to form a variety of different compounds. Plants and animals are made up of carbon compounds, and so are certain minerals. Carbon combines with oxygen to make a gas called carbon dioxide (CO₂).

Carbon cycle:

The movement and exchange of carbon through living organisms, the ocean, the atmosphere, rocks and minerals, and other parts of the Earth. Carbon moves from one place to another through various chemical, physical, geological, and biological processes.

Carbon dioxide:

A colorless, odorless greenhouse gas. It is produced naturally when dead animals or plants decay, and it is used by plants during photosynthesis. People are adding carbon dioxide into the atmosphere, mostly by burning fossil fuels such as coal, oil, and natural gas. This extra carbon dioxide is the main cause of climate change.

Greenhouse gas:

Also sometimes known as “heat-trapping gases,” greenhouse gases are natural or manmade gases that trap heat in the atmosphere and contribute to the greenhouse effect. Greenhouse gases include water vapor, carbon dioxide, methane, nitrous oxide, and fluorinated gases.

Parts per million (ppm):

A unit of measurement that can be used to describe the concentration of a particular substance within air, water, soil, or some other medium. For example, the concentration of carbon dioxide in the Earth's atmosphere is almost 400 parts per million, which means 1 million liters of air would contain about 400 liters of carbon dioxide.

Photosynthesis:

The process by which green plants use sunlight, water, and carbon dioxide to make food and other substances that they use to grow. In the process, plants release oxygen into the air.



CARBON THROUGH THE SEASONS

Mauna Loa on the World Map



Scientists monitor the amount of CO₂ and other gases in the atmosphere at stations such as the Mauna Loa Observatory in Hawaii. The Mauna Loa Observatory is one of the sites that have helped scientists determine that CO₂ levels in the atmosphere have increased significantly in recent decades and that these levels are continuing to rise at a rapid rate.

Image source: Carbon Dioxide Information Analysis Center: <http://cdiac.ornl.gov/>.


INSTRUCTIONS

Part 1: Plotting Monthly Atmospheric CO₂ Data

1. Tell the students that they will be learning about the carbon cycle by looking at monthly atmospheric CO₂ concentration data from 1959 to 2011. Explain that these data come from the Mauna Loa Observatory in Hawaii, and show students where the observatory is located on a map. Explain that because CO₂ spreads throughout the world's atmosphere, measurements from Mauna Loa are actually representative of global average CO₂ levels.
2. Show the class a short video about the carbon cycle and how people are changing this natural cycle at "Learn the Basics: Today's Climate Change" on EPA's *A Student's Guide to Global Climate Change* website (<http://www.epa.gov/climatechange/students/basics/today/carbon-dioxide.html>). You can also show a diagram of the carbon cycle (see box below), or demonstrate how the carbon cycle works by drawing it on the chalkboard.

CARBON THROUGH THE SEASONS

The Carbon Cycle



Adapted from: CO₂logic: <http://www.co2logic.com/home.aspx/html/Images/carbon%20cycle.jpg>.

- Discuss the processes of respiration and photosynthesis in plants and how these processes influence the amount of CO₂ in the atmosphere during the course of a year.
- Break the students into groups of approximately five. Assign each group a set of data from the “Mauna Loa Observatory Data” tables (each group will get 10 to 12 years of data), and pass out one copy of the assigned data per student. Also provide each group with a sheet of graph paper and a copy of the “Mauna Loa Worksheet.”
- Discuss what parts per million (ppm) means.
[Answer: Parts per million or ppm is a unit of measurement that can be used to describe the concentration of a particular substance within air, water, soil, or some other medium. For example, the concentration of carbon dioxide in the Earth's atmosphere is almost 400 parts per million, which means 1 million liters of air would contain about 400 liters of carbon dioxide.]

CARBON THROUGH THE SEASONS

6. Tell students that each group will be graphing their data on the graph paper. Explain that students in each group should work as a team by taking turns to plot data points on the same graph paper. Ask students to follow the “Instructions for Plotting the Graph” on the “Mauna Loa Data Worksheet” to plot their data.
7. When students have finished plotting their graphs, hand out a copy of the “Mauna Loa Monthly Average CO₂ Concentration, 1958 to 2011” graph to each group. Tell students that this is how the entire Mauna Loa Observatory data series looks when it is plotted in one graph.
8. Discuss the following questions in class:
 - What patterns do you notice in your graph? (Ask students to keep these patterns in mind as you ask them additional questions.)
[Answer: An increase in CO₂, followed by a decrease in CO₂, creating a repeating pattern of peaks and troughs much like a wave. Another pattern is an increase in heights of both peaks and troughs over time.]
 - What does the increase in the height of the peaks and troughs mean?
[Answer: The overall amount of CO₂ in the atmosphere is increasing.]
 - According to your data, during what month and during what season are the CO₂ concentrations highest? Lowest?
[Answer: highest concentrations in April and May (spring), lowest in August and September (early fall)]
 - Explain how the seasonal changes of CO₂ concentration in the atmosphere and the growing season for plants and are related?
[Answer: CO₂ in the atmosphere decreases during the growing season and increases during the rest of the year, which leads to maximum buildup in April and May before photosynthesis begins to take over again. Photosynthesis, in which plants take up CO₂ from the atmosphere and release oxygen, dominates during the growing season (the warmer part of the year). Respiration, by which plants and animals take up oxygen and release CO₂, occurs all the time but dominates during the colder months of the year.]
 - Are the seasons the same in the Northern and Southern Hemispheres?
[Answer: No, the seasons in the Northern and Southern Hemispheres are opposite.]
 - How does this difference affect CO₂ concentrations in the atmosphere?
[Answer: While CO₂ concentrations in the atmosphere are increasing in the Northern Hemisphere, CO₂ concentrations are decreasing in the Southern Hemisphere, and vice versa.]
 - There is more land in the Northern Hemisphere than in the Southern Hemisphere. How might this difference affect CO₂ concentrations in the atmosphere?
[Answer: The carbon cycle is more pronounced in the Northern Hemisphere (which has relatively more land mass and terrestrial vegetation) than in the Southern Hemisphere (which is more dominated by oceans).]
 - Earlier, you noticed that your line graph has a repeating pattern. Explain.
[Answer: The variations within each year are the result of the annual cycles of photosynthesis and respiration.]



CARBON THROUGH THE SEASONS

Part 2: Calculating Annual Average CO₂ Concentrations

[This portion of the lesson can be done in another class period or assigned as homework. It can be done as a group or individual exercise.]

1. Have each group of students complete the “Annual Average CO₂ Concentrations” chart on the “Mauna Loa Data Worksheet” for their set of data. Discuss the following questions:
 - Is there a particular pattern in the change in annual average CO₂ concentration for your set of years?
[Answer: Yes, CO₂ concentration increases every year. (Each group's dataset shows this same pattern.)]
 - Ask the students what this pattern means.
[Answer: It means more CO₂ was added to the atmosphere.]
2. Tell students that people are changing this Earth's natural carbon balance by adding more CO₂ to the atmosphere when we burn fossil fuels (such as coal, oil, and natural gas)—whether it's to drive our cars, power our homes, or make products. This extra CO₂ is being added to the atmosphere faster than natural processes can remove it, causing the atmosphere to trap more heat and causing the Earth's average temperature to rise.

EXTENSION

There are many monitoring stations like Mauna Loa Observatory that collect data about atmospheric CO₂ levels. Some scientists even use ice cores and tree rings to learn about the amount of CO₂ in the atmosphere hundreds of thousands of years ago.

- Tell students that the CO₂ levels in the past hundred years are abnormally high compared with natural historical levels, and that these levels are continuing to increase at an unprecedented rate.
- Explain that scientists can compare the amount of CO₂ in the atmosphere with the amount trapped in ancient ice cores, which show that the atmosphere had less CO₂ in the past.
- Show this three-minute YouTube video about the history of global atmospheric CO₂. (http://www.youtube.com/watch?feature=player_embedded&v=bbgUE04Y-Xg). The video starts with an animated graph of seasonal CO₂ variations observed at stations around the world from 1979 to 2011, then looks at patterns back to 800,000 years ago.



CARBON THROUGH THE SEASONS

MAUNA LOA OBSERVATORY DATA

Mauna Loa CO₂ Monthly Average Concentrations, 1959 to 1969 (ppm)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1959	315.62	316.38	316.71	317.72	318.29	318.15	316.54	314.80	313.84	313.26	314.80	315.58
1960	316.43	316.97	317.58	319.02	320.03	319.59	318.18	315.91	314.16	313.83	315.00	316.19
1961	316.93	317.70	318.54	319.48	320.58	319.77	318.57	316.79	314.80	315.38	316.10	317.01
1962	317.94	318.56	319.68	320.63	321.01	320.55	319.58	317.40	316.26	315.42	316.69	317.69
1963	318.74	319.08	319.86	321.39	322.25	321.47	319.74	317.77	316.21	315.99	317.12	318.31
1964	319.57	320.07	320.73	321.77	322.25	321.89	320.44	318.70	316.70	316.79	317.79	318.71
1965	319.44	320.44	320.89	322.13	322.16	321.87	321.39	318.81	317.81	317.30	318.87	319.42
1966	320.62	321.59	322.39	323.87	324.01	323.75	322.39	320.37	318.64	318.10	319.79	321.08
1967	322.07	322.50	323.04	324.42	325.00	324.09	322.55	320.92	319.31	319.31	320.72	321.96
1968	322.57	323.15	323.89	325.02	325.57	325.36	324.14	322.03	320.41	320.25	321.31	322.84
1969	324.00	324.42	325.64	326.66	327.34	326.76	325.88	323.67	322.38	321.78	322.85	324.11

Data source: National Oceanic and Atmospheric Administration (NOAA):
<http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>. Accessed August 2012.

Mauna Loa CO₂ Monthly Average Concentrations, 1970 to 1979 (ppm)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1970	325.03	325.99	326.87	328.13	328.07	327.66	326.35	324.69	323.10	323.16	323.98	325.13
1971	326.17	326.68	327.18	327.78	328.92	328.57	327.34	325.46	323.36	323.57	324.80	326.01
1972	326.77	327.63	327.75	329.72	330.07	329.09	328.05	326.32	324.93	325.06	326.50	327.55
1973	328.54	329.56	330.30	331.50	332.48	332.07	330.87	329.31	327.51	327.18	328.16	328.64
1974	329.35	330.71	331.48	332.65	333.19	332.12	330.99	329.17	327.41	327.21	328.34	329.50
1975	330.68	331.41	331.85	333.29	333.91	333.40	331.74	329.88	328.57	328.36	329.33	330.59
1976	331.66	332.75	333.46	334.78	334.78	334.06	332.95	330.64	328.96	328.77	330.18	331.65
1977	332.69	333.23	334.97	336.03	336.82	336.10	334.79	332.53	331.19	331.21	332.35	333.47
1978	335.10	335.26	336.61	337.77	338.01	337.98	336.48	334.37	332.33	332.41	333.76	334.83
1979	336.21	336.65	338.13	338.94	339.00	339.20	337.60	335.56	333.93	334.12	335.26	336.78

Data source: National Oceanic and Atmospheric Administration (NOAA):
<http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>. Accessed August 2012.



CARBON THROUGH THE SEASONS

MAUNA LOA OBSERVATORY DATA

Mauna Loa CO₂ Monthly Average Concentrations, 1980 to 1989 (ppm)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1980	337.80	338.28	340.04	340.86	341.47	341.26	339.34	337.45	336.10	336.05	337.21	338.29
1981	339.36	340.51	341.57	342.56	343.01	342.49	340.68	338.49	336.92	337.12	338.59	339.90
1982	340.92	341.69	342.85	343.92	344.67	343.78	342.23	340.11	338.32	338.39	339.48	340.88
1983	341.64	342.87	343.59	345.25	345.96	345.52	344.15	342.25	340.17	340.30	341.53	343.07
1984	344.05	344.77	345.46	346.77	347.55	346.98	345.55	343.20	341.35	341.68	343.06	344.54
1985	345.25	346.06	347.66	348.20	348.92	348.40	346.66	344.85	343.20	343.08	344.40	345.82
1986	346.54	347.13	348.05	349.77	350.53	349.90	348.11	346.09	345.01	344.47	345.86	347.15
1987	348.38	348.70	349.72	351.32	352.14	351.61	349.91	347.84	346.52	346.65	347.95	349.18
1988	350.38	351.68	352.24	353.66	354.18	353.68	352.58	350.66	349.03	349.08	350.15	351.44
1989	352.89	353.24	353.80	355.59	355.89	355.30	353.98	351.53	350.02	350.29	351.44	352.84

Data source: National Oceanic and Atmospheric Administration (NOAA):
<http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>. Accessed August 2012.

Mauna Loa CO₂ Monthly Average Concentrations, 1990 to 1999 (ppm)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1990	353.79	354.88	355.65	356.27	357.29	356.32	354.89	352.89	351.28	351.59	353.05	354.27
1991	354.87	355.68	357.06	358.51	359.09	358.10	356.12	353.89	352.30	352.32	353.79	355.07
1992	356.17	356.93	357.82	359.00	359.55	359.32	356.85	354.91	352.93	353.31	354.27	355.53
1993	356.86	357.27	358.36	359.27	360.19	359.52	357.42	355.46	354.10	354.12	355.40	356.84
1994	358.22	358.98	359.91	361.32	361.68	360.80	359.39	357.42	355.63	356.09	357.56	358.87
1995	359.87	360.79	361.77	363.23	363.77	363.22	361.70	359.11	358.11	357.97	359.40	360.61
1996	362.04	363.17	364.17	364.51	365.16	364.93	363.53	361.38	359.60	359.54	360.84	362.18
1997	363.04	364.09	364.47	366.25	366.69	365.59	364.34	362.20	360.31	360.71	362.44	364.33
1998	365.18	365.98	367.13	368.61	369.49	368.95	367.74	365.79	364.01	364.35	365.52	367.08
1999	368.12	368.98	369.60	370.96	370.77	370.33	369.28	366.86	364.94	365.35	366.68	368.04

Data source: National Oceanic and Atmospheric Administration (NOAA):
<http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>. Accessed August 2012.



CARBON THROUGH THE SEASONS

MAUNA LOA OBSERVATORY DATA

Mauna Loa CO₂ Monthly Average Concentrations, 2000 to 2011 (ppm)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
2000	369.25	369.50	370.56	371.82	371.51	371.71	369.85	368.20	366.91	366.99	368.33	369.67
2001	370.52	371.49	372.53	373.37	373.82	373.18	371.57	369.63	368.16	368.42	369.69	371.18
2002	372.45	373.14	373.93	375.00	375.65	375.50	374.00	371.83	370.66	370.51	372.20	373.71
2003	374.87	375.62	376.48	377.74	378.50	378.18	376.72	374.31	373.20	373.10	374.64	375.93
2004	377.00	377.87	378.73	380.41	380.63	379.56	377.61	376.15	374.11	374.44	375.93	377.45
2005	378.47	379.76	381.14	382.20	382.47	382.20	380.78	378.73	376.66	376.98	378.29	379.92
2006	381.35	382.16	382.66	384.73	384.98	384.09	382.38	380.45	378.92	379.16	380.18	381.79
2007	382.93	383.81	384.56	386.40	386.58	386.05	384.49	382.00	380.90	381.14	382.42	383.89
2008	385.44	385.73	385.97	387.16	388.50	387.88	386.42	384.15	383.09	382.99	384.13	385.56
2009	386.93	387.42	388.77	389.46	390.19	389.44	387.91	385.92	384.79	384.39	386.00	387.27
2010	388.45	389.82	391.08	392.46	392.95	392.03	390.13	388.15	386.80	387.18	388.59	389.68
2011	391.19	391.76	392.40	393.28	394.16	393.68	392.39	390.08	389.00	388.92	390.20	391.80

Data source: National Oceanic and Atmospheric Administration (NOAA):
<http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>. Accessed August 2012.



CARBON THROUGH THE SEASONS

MAUNA LOA DATA WORKSHEET

NAME: _____ DATE: _____

Instructions for Plotting the Graph

1. On the graph paper, label the x-axis as "Year and Month." Begin numbering from left to right on the x-axis from 1 to 12 at intervals of 1. Each point represents a month of the year: 1 is January, 2 is February, and so on. Label this first group of 12 points (or months) below the x-axis as the first year of your assigned data. Continue numbering points on the x-axis in sets of 12 months for the next years.
2. On the y-axis (on the left-hand side of the paper), number the parts per million for CO₂ concentrations. Begin with 300 at the lower end, and number up to 400 ppm, counting by intervals of 10. Label the axis.
3. Using information from the "Mauna Loa Observatory Data," table, plot the points corresponding to each monthly average atmospheric CO₂ concentration for the years assigned to your group.

Instructions for Filling Out the Changes in Annual Average CO₂ Concentrations

Fill in the years of your individual dataset in the chart below. Calculate 1) the average (mean) CO₂ concentration (ppm) for each of the years in your dataset, 2) the amount of change (ppm) from the previous year, and 3) the direction of change (whether it has increased, decreased, or remained unchanged).


Annual Average CO₂ Concentrations

Year	Annual average	Difference from previous year	Direction



CARBON THROUGH THE SEASONS

Mauna Loa Monthly Average CO₂ Concentration, 1958 to 2011



The Mauna Loa Observatory is one of the sites that have helped scientists determine that CO₂ levels in the atmosphere have increased significantly in recent decades and that these levels are continuing to rise at a rapid rate. CO₂ stays in the atmosphere for long enough that it is able to spread fairly evenly around the world, so even measurements from a single site (like Mauna Loa) can be representative of global average CO₂ concentrations.

Image source: National Oceanic and Atmospheric Administration (NOAA):
http://www.esrl.noaa.gov/gmd/webdata/ccgg/trends/co2_data_mlo.png. Accessed September 2012.

For the latest version, please see the graph in the “Full Mauna Loa CO₂ Record” section of NOAA’s page at <http://www.esrl.noaa.gov/gmd/ccgg/trends/mlo.html>.

CARBON THROUGH THE SEASONS

ANNUAL AVERAGE CO₂ CONCENTRATIONS—ANSWER KEY

Year	Annual average	Difference from previous year	Direction
1959	315.97		
1960	316.91	0.93	Increase
1961	317.64	0.73	Increase
1962	318.45	0.81	Increase
1963	318.99	0.54	Increase
1964	319.62	0.62	Increase
1965	320.04	0.43	Increase
1966	321.38	1.34	Increase
1967	322.16	0.77	Increase
1968	323.05	0.89	Increase
1969	324.62	1.58	Increase
1970	325.68	1.06	Increase
1971	326.32	0.64	Increase
1972	327.45	1.13	Increase
1973	329.68	2.22	Increase
1974	330.18	0.50	Increase
1975	331.08	0.91	Increase
1976	332.05	0.97	Increase
1977	333.78	1.73	Increase
1978	335.41	1.63	Increase
1979	336.78	1.37	Increase
1980	338.68	1.90	Increase
1981	340.10	1.42	Increase
1982	341.44	1.34	Increase
1983	343.03	1.59	Increase
1984	344.58	1.55	Increase
1985	346.04	1.46	Increase
1986	347.38	1.34	Increase
1987	349.16	1.78	Increase
1988	351.56	2.40	Increase
1989	353.07	1.50	Increase
1990	354.35	1.28	Increase

(CONTINUED)



CARBON THROUGH THE SEASONS

ANNUAL AVERAGE CO₂ CONCENTRATIONS—ANSWER KEY (CONTINUED)

Year	Annual average	Difference from previous year	Direction
1991	355.57	1.22	Increase
1992	356.38	0.82	Increase
1993	357.07	0.69	Increase
1994	358.82	1.76	Increase
1995	360.80	1.97	Increase
1996	362.59	1.79	Increase
1997	363.71	1.12	Increase
1998	366.65	2.95	Increase
1999	368.33	1.67	Increase
2000	369.53	1.20	Increase
2001	371.13	1.61	Increase
2002	373.22	2.08	Increase
2003	375.77	2.56	Increase
2004	377.49	1.72	Increase
2005	379.80	2.31	Increase
2006	381.90	2.10	Increase
2007	383.76	1.86	Increase
2008	385.59	1.82	Increase
2009	387.37	1.79	Increase
2010	389.78	2.40	Increase
2011	391.57	1.79	Increase



Hudson River Estuary Climate Change Lesson Project

www.nyseagrant.org
<http://www.dec.ny.gov/lands/>



New York State
Water Resources Institute



Cornell University